arXiv:math/0410490v1 [math.CA] 22 Oct 2004

Potpourri, 7

Stephen William Semmes
Rice University
Houston, Texas

Let V be a finite-dimensional real or complex vector space, with positive dimension, and let $\|v\|_V$ be a norm on V. Thus $\|v\|_V$ is a nonnegative real number for all $v \in V$, $\|v\|_V = 0$ if and only if $v = 0$, $\|\alpha\, v\|_V = |\alpha|\,\|v\|_V$ for all real or complex numbers α, as appropriate, and all $v \in V$, and $\|v + w\|_V \le \|v\|_V + \|w\|_V$ for all $v, w \in V$.

Associated to this norm we get a metric $\|v - w\|_V$ on V. Since V has finite dimension, it is linearly isomorphic to $\mathbf{R}^n$ or $\mathbf{C}^n$, where n is the dimension of V, according to whether V is a real or complex vector space. The topology on V determined by the metric $\|v - w\|_V$ corresponds exactly to the usual topology on $\mathbf{R}^n$ or $\mathbf{C}^n$ with respect to this isomorphism, by standard arguments.

Let λ be a linear functional on V, which is to say a linear mapping from V into the real or complex numbers, as appropriate. Using the identification with a Euclidean space, one can see that there is a nonnegative real number k such that

$$|\lambda(v)| \le k\,\|v\|_V \tag{1}$$

for all $v \in V$.

Put

$$\|\lambda\|_{V^*} = \sup\{|\lambda(v)| : v \in V,\ \|v\|_V \le 1\}. \tag{2}$$

The supremum is finite by the remarks in the preceding paragraph, and $\|\lambda\|_{V^*}$ can also be described as the smallest nonnegative real number k such that $|\lambda(v)|$ is less than or equal to k times $\|v\|_V$ for all $v \in V$. The space of all linear functionals on V is denoted V^* and is a vector space with the same dimension as V, and $\|\lambda\|_{V^*}$ defines a norm on V^* called the dual norm associated to $\|v\|_V$.

If v is any vector in V, then a famous theorem implies that there is a linear functional λ on V such that $\|\lambda\|_{V^*} = 1$ and

$$\lambda(v) = \|v\|_V. \tag{3}$$

It follows that

$$\|v\|_V = \max\{|\lambda(v)| : \lambda \in V^*,\ \|\lambda\|_{V^*} \le 1\} \tag{4}$$

for all $v \in V$.

Let Σ denote the unit sphere in V^* with respect to the dual norm $\|\lambda\|_{V^*}$, i.e.,

$$\Sigma = \{\lambda \in V^* : \|\lambda\|_{V^*} = 1\}. \tag{5}$$

This is a compact subset of V^* with respect to the topology induced by the dual norm, which is the same as a topology induced from the standard Euclidean topology using an isomorphism with Euclidean space. Let us write $C(\Sigma)$ for the vector space of continuous real or complex-valued functions on Σ, using the same scalars as for V.

For $\phi \in C(\Sigma)$ put

$$\|\phi\| = \sup\{|\phi(\lambda)| : \lambda \in \Sigma\}, \tag{6}$$

where in fact the maximum is attained by standard results in analysis. One can check that this defines a norm on $C(\Sigma)$, called the supremum norm.

For each $v \in V$ we get a function $\phi_v \in C(\Sigma)$ given by

$$\phi_v(\lambda) = \lambda(v). \tag{7}$$

The correspondence $v \mapsto \phi_v$ defines a linear mapping from V into $C(V)$ such that

$$\|\phi_v\| = \|v\|_V \tag{8}$$

for all $v \in V$.

Let E be a finite nonempty set, and let $\mathcal{F}(E, V)$ denote the vector space of V-valued functions on E. If f is a real or complex-valued function on E, put

$$\|f\|_\infty = \max\{|f(x)| : x \in E\}, \tag{9}$$

which defines a norm on the vector space of real or complex-valued functions on E. Similarly, if f is a V-valued function on E, then put

$$\|f\|_{\infty,V} = \max\{\|f(x)\|_V : x \in E\}, \tag{10}$$

which defines a norm on $\mathcal{F}(E, V)$.

Suppose that $t(x, y)$ is a real or complex-valued function on the Cartesian product $E \times E$. Thus we get a linear mapping T from the vector space of real or complex-valued functions on E to itself, as appropriate, given by

$$T(f)(x) = \sum_{y \in E} t(x, y)\, f(y). \tag{11}$$

Let a be a nonnegative real number such that

$$\|T(f)\|_\infty \le a\, \|f\|_\infty \tag{12}$$

for all real or complex-valued functions f on E. This is equivalent to the condition that

$$\sum_{y \in E} |t(x, y)| \le a \tag{13}$$

for all $x \in E_2$.

Let us write T_V for the version of T acting on V-valued functions on E, assuming that V is real or complex according to whether $t(x, y)$ is real or complex-valued. Namely, for a V-valued function f on E, $T_V(f)(x)$ is defined by the same formula as in the scalar case. One can check that

$$\|T_V(f)\|_{\infty,V} \le a\, \|f\|_{\infty,V} \tag{14}$$

for all V-valued functions f on E, using the condition on $|t(x,y)|$ mentioned in the previous paragraph. Alternatively, one can consider the linear operator $T_{C(\Sigma)}$ which is the version of T on $C(\Sigma)$-valued functions on E, which basically acts pointwise on Σ, and derive the analogous inequality using the supremum norm on $C(\Sigma)$ from the inequality for scalar-valued functions. The inequality for V-valued functions then follows by employing the isometric embedding of V in $C(\Sigma)$.

If f is a real or complex-valued function on E, then put

$$\|f\|_1 = \sum_{x \in E} |f(x)|, \tag{15}$$

and more generally if f is a V-valued function on E put

$$\|f\|_{1,V} = \sum_{x \in E} \|f(x)\|_V. \tag{16}$$

These define norms on the vector spaces of scalar-valued and V-valued functions on E.

Suppose that $t(x,y)$, T are as before, and that b is a nonnegative real number such that

$$\|T(f)\|_1 \le b\,\|f\|_1 \tag{17}$$

for all scalar-valued functions f on E. This is equivalent to the condition that

$$\sum_{x \in E} |t(x,y)| \le b \tag{18}$$

for all $x \in E$. In this event we have that

$$\|T(f)\|_{1,v} \le b\,\|f\|_{1,V} \tag{19}$$

for all V-valued functions f on E too.

Fix a real number p, $1 < p < \infty$, and put

$$\|f\|_p = \biggl(\sum_{x \in E} |f(x)|^p\biggr)^{1/p} \tag{20}$$

for scalar-valued functions on E and

$$\|f\|_{p,V} = \biggl(\sum_{x \in E} \|f(x)\|_V^p\biggr)^{1/p} \tag{21}$$

for V-valued functions on E. Again these define norms on the vector spaces of scalar and V-valued functions on E.

Let us take for V the vector space of real or complex-valued functions on E, as appropriate. A V-valued function on E is then basically the same as a real or complex-valued function $f(x,z)$ on $E \times E$. The linear transformation T_V on

V-valued functions on E can be described explicitly by saying that if $f(w, z)$ is a real or complex-valued function on $E \times E$, as appropriate, then $T_V(f)$ is the function on $E \times E$ given by

$$T_V(f)(x, z) = \sum_{y \in E} t(x, y)\, f(y, z). \tag{22}$$

Let us use the norm $\| \cdot \|_1$ on functions on E as our norm on V. If $1 \le p < \infty$ and $f(x, z)$ is a function on $E \times E$, which is equivalently a V-valued function on E, then

$$\|f\|_{p,V} = \biggl(\sum_{x \in E} \biggl(\sum_{z \in E} |f(x, z)| \biggr)^p \biggr)^{1/p}. \tag{23}$$

Suppose that $h(x)$ is a real or complex-valued function on E, as appropriate. Consider the function $H(x, z)$ on $E \times E$ given by $H(x, z) = h(x)$ when $x = z$, $H(x, z) = 0$ when $x \ne z$. Thus we have that $\|H\|_{p,V} = \|h\|_p$ for all p, $1 \le p < \infty$.

Fix p, $1 < p < \infty$, and suppose that B_p is a nonnegative real number such that

$$\|T_V(f)\|_{p,V} \le B_p \, \|f\|_{p,V} \tag{24}$$

for all V-valued functions on E. If h is a scalar-valued function on E and H is the associated function on $E \times E$ described in the previous paragraph, then $T_V(H)$ is the function on $E \times E$ given by $t(x, z)\, h(z)$. Hence

$$\biggl(\sum_{x \in E} \biggl(\sum_{z \in E} |t(x, z)| \, |h(z)| \biggr)^p \biggr)^{1/p} \le B_p \biggl(\sum_{w \in E} |h(w)|^p \biggr)^{1/p} \tag{25}$$

for all real or complex-valued functions h on E.

References

[Bea1] R. Beals, *Topics in Operator Theory*, The University of Chicago Press, 1971.

[Bea2] R. Beals, *Advanced Mathematical Analysis: Periodic Functions and Distributions, Complex Analysis, Laplace Transform and Applications*, Springer-Verlag, 1973.

[BenS] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, 1988.

[BenL] Y. Benyamini and J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, Volume 1, Colloquium Publications **48**, American Mathematical Society, 2000.

[BerL] J. Bergh and J. Löfström, *Interpolation Spaces*, Springer-Verlag, 1976.

[Boc] S. Bochner, *Lectures on Fourier Integrals*, with an author's supplement on monotonic functions, Stieltjes integrals, and harmonic analysis, translated from the 1932 original by M. Tenenbaum and H. Pollard, Annals of Mathematics Studies **42**, Princeton University Press, 1959.

[BocC] S. Bochner and K. Chandrasekharan, *Fourier Transforms*, Annals of Mathematics Studies **19**, Princeton University Press, 1949.

[Duo] J. Duoandikoetxea, *Fourier Analysis*, translated and revised by D. Cruz-Uribe, SFO, American Mathematical Society, 2001.

[GarR] J. García-Cuerva and J. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland, 1985.

[Gol] R. Goldberg, *Methods of Real Analysis*, 2nd edition, Wiley, 1976.

[HewS] E. Hewitt and K. Stromberg, *Real and Abstract Analysis*, Springer-Verlag, 1975.

[JohL] W. Johnson and J. Lindenstrauss, editors, *Handbook of the Geometry of Banach Spaces*, Volumes I and II, North-Holland, 2001 and 2003.

[Jon] F. Jones, *Lebesgue Integration on Euclidean Space*, Jones and Bartlett, 1993.

[Jou] J.-L. Journé, *Calderón–Zygmund Operators, Pseudodifferential Operators, and the Cauchy Integral of Calderón*, Lecture Notes in Mathematics **994**, Springer-Verlag, 1993.

[KalPR] N. Kalton, N. Peck, and J. Roberts, *An F-Space Sampler*, Cambridge University Press, 1984.

[Kat] Y. Katznelson, *An Introduction to Harmonic Analysis*, 3rd edition, Cambridge University Press, 2004.

[Kra1] S. Krantz, *Real Analysis and Foundations*, CRC Press, 1991.

[Kra2] S. Krantz, *A Panorama of Harmonic Analysis*, Carus Mathematical Monographs **27**, Mathematical Association of America, 1999.

[LinT] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I, Sequence Spaces, II, Function Spaces*, Springer-Verlag, 1977, 1979.

[McC] C. McCarthy, c_p, Israel Journal of Mathematics **5** (1967), 249–271.

[Mil] M. Milman, *Extrapolation and Optimal Decompositions with Applications to Analysis*, Lecture Notes in Mathematics **1580**, Springer-Verlag, 1994.

[MilS] V. Milman and G. Schechtman, *Asymptotic Theory of Finite-Dimensional Normed Spaces*, with an appendix by M. Gromov, Lecture Notes in Mathematics **1200**, Springer-Verlag, 1986.

[Pelc] A. Pełczyński, *Banach Spaces of Analytic Functions and Absolutely Summing Operators*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics **30**, American Mathematical Society, 1977.

[Pell] V. Peller, *Hankel Operators and their Applications*, Springer-Verlag, 2003.

[Pis1] G. Pisier, *Factorization of Linear Operators and Geometry of Banach Spaces*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics **60**, American Mathematical Society, 1986.

[Pis2] G. Pisier, *The Volume of Convex Bodies and Banach Space Geometry*, Cambridge University Press, 1989.

[Rud1] W. Rudin, *Principles of Mathematical Analysis*, 3rd edition, McGraw-Hill, 1976.

[Rud2] W. Rudin, *Real and Complex Analysis*, 3rd edition, McGraw-Hill, 1987.

[Rud3] W. Rudin, *Functional Analysis*, 2nd edition, McGraw-Hill, 1991.

[Sad] C. Sadosky, *Interpolation of Operators and Singular Integrals: An Introduction to Harmonic Analysis*, Marcel Dekker, 1979.

[Sim] B. Simon, *Trace Ideals and their Applications*, Cambrdige University Press, 1979.

[Ste1] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.

[Ste2] E. Stein, *The development of square functions in the work of A. Zygmund*, Bulletin of the American Mathematical Society (New Series) **7** (1982), 359–376.

[Ste3] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.

[SteS1] E. Stein and R. Shakarchi, *Fourier Analysis: An Introduction*, Princeton University Press, 2003.

[SteS2] E. Stein and R. Shakarchi, *Complex Analysis*, Princeton University Press, 2003.

[SteS3] E. Stein and R. Shakarchi, *Real Analysis*, Princeton University Press, to appear.

[SteS4] E. Stein and R. Shakarchi, *Probability Theory*, Princeton University Press, to appear.

[SteW1] E. Stein and G. Weiss, *Interpolation of operators with change of measures*, Transactions of the American Mathematical Society **87** (1958), 159–172.

[SteW2] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.

[StrT] J.-O. Strömberg and A. Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Mathematics **1381**, Springer-Verlag, 1989.

[Tor] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Dover, 2004.

[Tre] F. Treves, *Topological Vector Spaces, Distributions, and Kernels*, Academic Press, 1967.

[WheZ] R. Wheeden and A. Zygmund, *Measure and Integral: An Introduction to Real Analysis*, Marcel Dekker, 1977.

[Woj] P. Wojtaszczyk, *Banach Spaces for Analysts*, Cambridge University Press, 1991.

[Zem] A. Zemanian, *Distribution Theory and Transform Analysis: An Introduction to Generalized Functions, with Applications*, 2nd edition, Dover, 1987.

[Zyg] A. Zygmund, *Trigonometric Series*, Volumes I and II, 3rd edition, Cambridge University Press, 2002.